


From High-Energy C₇H₂ Isomers with A Planar Tetracoordinate Carbon Atom to An Experimentally Known Carbene

Krishnan Thirumoorthy,[†] Amir Karton,^{*,‡} and Venkatesan S. Thimmakondur^{*,§}[†]Department of Chemistry, School of Advanced Sciences, Vellore Institute of Technology, Vellore 632 014, Tamil Nadu, India[‡]School of Molecular Sciences, The University of Western Australia, Perth, Western Australia 6009, Australia[§]Department of Chemistry, Birla Institute of Technology and Science, Pilani, K K Birla Goa Campus, Goa 403 726, India

S Supporting Information

ABSTRACT: In this work, we use high-level ab initio procedures to show that the high-energy isomers of C₇H₂ with a planar tetracoordinate carbon (ptC) atom serve as reactive intermediate leading to the formation of an experimentally known ring-chain carbene, 1-(buta-1,3-diynyl)-cyclopropenyliene (**2**). Among the experimentally known isomers of C₇H₂, the latter is the only low-lying ring-chain carbene identified by Fourier-transform microwave spectroscopy. Here we investigate the ring-opening pathways of C–C single bonds connected to the ptC atom in three different C₇H₂ isomers using coupled-cluster and density functional theory methods. These three isomers [ptC1 (C_{2v}; \tilde{X}^1A_1), ptC2 (C_s; \tilde{X}^1A'), and ptC3 (C_s; \tilde{X}^1A')] are found to be local minima on the C₇H₂ potential-energy surface at both CCSD(T)/cc-pVTZ and B3LYP/6-311+G(d,p) levels of theory. The transition states and minimum-energy pathways connecting the reactants (ptC isomers) and the products have been found via intrinsic reaction coordinate calculations at the B3LYP/6-311+G(d,p) level of theory. The high-energy ptC isomers (ptC2 and ptC3) lead to the formation of **2**, while the low-energy ptC isomer, ptC1, rearranges to a bicyclic carbene, bicyclo[4.1.0]hepta-4,6-diene-2-yne-7-ylidene (**6**). In the latter, we note that both the reactant and the product are yet to be identified in the laboratory. Relative energies, activation energies, reaction energies, and nucleus independent chemical shift values have been calculated to access the thermodynamic and kinetic stabilities and the aromatic nature of these peculiar molecules. Rotational and centrifugal distortion constants have also been estimated for all ptC isomers, which may assist the efforts of microwave spectroscopists.



INTRODUCTION

In a given elemental composition, the limelight of hypothetical molecules may always remain poor unless and until they are identified in the laboratory. The energy difference between the low-energy isomers and the hypothetical molecules predicted to be local minima computationally is often an important factor in the acceptance or negligence of the latter. However, it is important to consider potential rearrangements or dissociations involving unidentified high-energy intermediates. To date, the concept of planar tetracoordinate carbon (ptC) has rather been seen as an odd feature as they can be considered as *anti van't Hoff-Le Bel* compounds.^{1–3} Nevertheless, this concept has gained some momentum experimentally^{4–8} though it was suggested nearly 50 years ago. First introduced by Monkhurst,⁹ the idea of how to stabilize a molecule containing a ptC atom was started by Hoffmann and co-workers.¹⁰ Although only a handful of molecules were identified in the laboratory,^{4–8} theoretically many molecules have been proposed to date. These include hydrocarbons stabilized by electronic effects^{11–17} as well as hydrocarbons stabilized by steric effects such as alkylplanes and hemispiroalkylplanes.^{18–22} Doped hydrocarbons or carbon clusters

with lithium,²³ beryllium/boron,^{24,25} boron,^{26–30} nitrogen,³¹ silicon,³² phosphorus,^{33,34} and late transition metals³⁵ with a ptC, ptB, ptN, ptSi, or ptP atom are also reported. It is worth to note here that planar penta-,^{36–43} hexa-,^{44–48} and hepta-^{49,50} coordinated carbon compounds are also actively pursued due to the potential implications of these molecules in making new materials.⁵¹

In this work, we explored the dissociation pathways of three different isomers of C₇H₂, which contain a ptC atom. ptC1, ptC2, and ptC3 (see Figure 1) can be named as 2,7-(didehydro)-tricyclo[4.1.0.0^{1,3}]hepta-2,4,6-triene, 2,5-(didehydro)-1-ethynylspiro[2.2]penta-1,4-diene, and 2,4-(didehydro)-1-ethynylspiro[2.2]penta-1,4-diene, respectively.⁵² The equilibrium geometry of ptC1 has been reported elsewhere more than a decade before.^{15,16} Nevertheless, it remains elusive in the laboratory to date though energetically lying close to 1,1-(diethynyl)propadienyliene (**11**) and the cumulene carbene isomer of C₇H₂, heptahexaenyliene (**9**), which are exper-

Received: September 9, 2018

Revised: October 25, 2018

Published: October 26, 2018




Figure 1. Various isomers of C_7H_2 considered in this and previous theoretical works.^{59,60} The relative energy differences were calculated at the W3lite-F12 level of theory.¹²³ ZPVEs are calculated at the CCSD(T)/cc-pVTZ level and scaled by a factor of 0.9868.⁹⁹ The experimentally detected isomers are marked with an asterisk symbol. Isomers 1 and 16 are triplets and all others are singlets.




Figure 2. Schematic reaction profile diagram for **ptC1** and its dissociation pathways connected to the **ptC** atom. The relative energy differences were calculated at the W1-F12 level of theory.

imentally known.^{53,54} The latter molecule and also 1-(buta-1,3-diyne)cyclopropenylidene (**2**) have been identified using Fourier transform microwave (FTMW) spectroscopy by McCarthy and co-workers.^{53,55} To the best of our knowledge, the important spectroscopic parameters such as rotational and centrifugal distortion constants have not been reported in other theoretical works to date,^{15,16,32} for **ptC1**, which may aid the detection of this peculiar molecule using FTMW spectroscopy in the laboratory.

The high-energy **ptC** isomers, **ptC2** and **ptC3**, have not been considered before theoretically.^{15,16,54,56–60} On the basis of CCSD(T)/cc-pVTZ//UB3LYP/6-311G(d,p) calculations, Sun et al. have proposed 113 isomers of C_7H_2 ;⁵⁸ however, **ptC** isomers were not considered in their study. To date, six isomers of C_7H_2 have been detected experimentally.^{53–55,61–63} Interest in C_7H_2 isomers largely stems from the fact that four carbene molecules of the lower homologous series [cyclopropenylidene ($c-C_3H_2$),⁶⁴ propadienylidene (C_3H_2),⁶⁵ butatrienylidene (C_4H_2),⁶⁶ and hexapentaenylidene (C_6H_2)⁶⁷]

Table 1. Dissociation Pathways of C_7H_2 Isomers with a ptC Atom and Their Corresponding Activation Energies (ΔE^\ddagger) and Reaction Energies (ΔE_r) Computed at the W1–F12 Level of Theory

isomer	dissociation pathway	ΔE^\ddagger (kcal mol ⁻¹)	ΔE_r (kcal mol ⁻¹)	NICS (1 Å) (ppm)	ΔE_0^c (kcal mol ⁻¹)
ptC1	A	11.02	-5.65 (6) ^a	-11.45	26.12
	B	32.96	(6) ^a		
ptC2	A	9.25	-46.83 (2) ^a	-7.72	52.53
	B	10.56	(2) ^a		
	C	6.11	3.27 (BC) ^b		
	D	10.21	-44.93 (3) ^a		
ptC3	A	7.45	-50.11 (2) ^a	-9.02	55.74
	B	7.76	(2) ^a		
	C	8.33	-48.21 (3) ^a		
	D	2.83	0.00 (BC) ^b		

^aThe dissociation pathway leading to the corresponding product is given in parentheses. The reaction path is confirmed by IRC calculations. ^bThis particular reaction path is not confirmed by IRC calculations. However, reoptimization from the last point of IRC calculation leads to this product. ^cW3lite-F12 relative energies with respect to isomer 1 at 0 K.




Figure 3. Schematic reaction profile diagram for ptC2 and its dissociation pathways connected to the ptC atom. The relative energy differences were calculated at the W1–F12 level of theory.

have been identified in interstellar sources. We also emphasize here that detection of these molecules in the laboratory^{64,68–70} is a crucial factor in the confirmation of several identified lines in the astronomical sources. The isotopologue of C_3H_2 , the doubly deuterated cyclopropenylidene ($c-C_3D_2$) has also recently been found in the interstellar sources.⁷¹ Although there is no evidence for the isomers of C_7H_2 in the spectral line surveys of IRC+10216⁷² and W51,⁷³ chemical models of Herbst and Leung⁷⁴ suggest that the abundance of C_7H_2 (unspecific to any particular isomer) in TMC-1 is comparable to HC_3N ^{75,76} and C_3N ,⁷⁷ which were observed in space. However, based on the available laboratory data of C_7H_2 isomers,^{53,55} McCarthy and co-workers had stated that the limits on the abundances of C_7H_2 isomers were not determined in the carbon-chain rich source of TMC-1 as their spectral line surveys have not been published yet. Here, our interest in ptC isomers of C_7H_2 emerge from the fact that their dissociation predominantly leads to low-energy isomers (2, 1,2-(diethynyl)cyclopropenylidene (3), and bicyclo[4.1.0]-

hepta-4,6-diene-2-yne-7-ylidene (6); see Table 1) depending upon the ptC isomer, and its corresponding C–C single bond connected to the ptC atom being broken. In particular, the major product obtained upon the dissociation (see Figures 3 and 4) of the high-energy ptC isomers (ptC2 and ptC3) is the experimentally known molecule of C_7H_2 , 2.⁵⁵ Therefore, one can unequivocally say that the latter serve as reactive intermediates within equilibrium thermodynamic considerations. It is also worth mentioning here that the production and identification of C_7H_2 isomers using FTMW spectroscopy involved an electrical discharge of 1% mixture of diacetylene in Ne with the supersonic nozzle of about 1000 V.^{55,78} The latter quantity certainly does not indicate the amount of energy being imparted on the precursor gases. Nevertheless, understanding the thermochemistry of C_7H_2 isomers on the high-energy regions of the potential energy surface (PES) is important for predicting potential rearrangements. While we do not discard the formation of C_7H_2 isomers by radical recombination of diacetylene subunits, the scope of our work




Figure 4. Schematic reaction profile diagram for **ptC3** and its dissociation pathways connected to the **ptC** atom. The relative energy differences were calculated at the W1–F12 level of theory.

in this paper is limited to equilibrium thermodynamic considerations.

In continuation of earlier theoretical^{15,16,54,56–58} and experimental studies on C_7H_2 isomers,^{53–55,61–63} in this work, we explored the possibilities of arriving at new isomers of C_7H_2 with a **ptC** atom. **ptC1** was initially reported as a local minimum on the C_7H_2 PES elsewhere at the B3LYP/6-311++G(2d,2p) level of theory.^{15,16} Nevertheless, our theoretical study differs from the earlier theoretical works as far as high-energy **ptC** isomers (**ptC2** and **ptC3**) and their dissociation are concerned. Our main motivation in this work is to determine the relative energies (ΔE_0), activation energies (ΔE^\ddagger), and also the reaction energies (ΔE_r) by studying their dissociation, which may give an idea of the thermodynamic versus kinetic stabilities of these peculiar molecules. We have also computed nucleus independent chemical shift (NICS) values^{79,80} for all the three **ptC** isomers of C_7H_2 to gauge the aromatic nature of these compounds. Optimal geometries and spectroscopic parameters, which are especially relevant to the FTMW spectroscopy are also collected. We believe that our efforts would assist the efforts of the experimentalists in the identification of these peculiar molecules in the future.

■ COMPUTATIONAL METHODS

The geometries of all C_7H_2 isomers with a **ptC** atom and their corresponding dissociative products were optimized using second-order Møller–Plesset perturbation theory,⁸¹ density functional theory (DFT), and also with coupled-cluster singles and doubles^{82,83} augmented with perturbative treatments of triple excitations (CCSD(T)) to incorporate a high-level treatment of electron correlation effects.^{84–86} For the MP2 and CCSD(T) calculations, the correlation-consistent polarized valence n zeta (cc-pVnZ) basis sets of Dunning's ($n = D$ and T) were employed.⁸⁷ For **ptC** isomers alone, we have also done geometry optimizations at the MP2/aug-cc-pVQZ level of theory to check consistencies in bond lengths and bond angles. The frozen-core approximation is utilized in all MP2

and CCSD(T) calculations. These calculations were done with the CFOUR⁸⁸ and MOLPRO^{89,90} program packages. All DFT calculations were done with the B3LYP hybrid-functional^{91,92} using the 6-311+G(d,p) basis set.^{93,94} All the transition states corresponding to the dissociation of C–C bonds connected to the **ptC** atom were obtained at the latter level. IRC calculations^{95,96} and NICS values^{79,80} were also estimated at the same level. NICS values reported here have been calculated at 1 Å above the plane of the **ptC** isomers. These calculations were done with the Gaussian 09 program.⁹⁷ We note that for all the stationary points obtained, harmonic vibrational frequencies were calculated by analytic calculation of second derivatives.⁹⁸ Zero-point vibrational energies (ZPVEs) calculated at the CCSD(T)/cc-pVTZ level of theory are scaled by a scaling factor of 0.9868 as recommended elsewhere.⁹⁹

In order to obtain reliable relative energies and reaction profiles, high-level benchmark data have been obtained using W1–F12 and W3lite-F12 theories.^{100–104} These theories represent layered extrapolations to the relativistic, all-electron CCSD(T)/CBS (W1–F12) and CCSDT(Q)/CBS (W3lite-F12) levels. W3lite-F12 theory can achieve benchmark accuracy for atomization reactions (i.e., it is associated with root-mean-square deviations, RMSDs, from accurate atomization energies of about $1 \text{ kJ mol}^{-1} = 0.24 \text{ kcal mol}^{-1}$). For example, the related W3–F12 theory is associated with an RMSD of $0.27 \text{ kcal mol}^{-1}$ for a set of 140 very accurate atomization energies obtained at the full configuration interaction (FCI) infinite basis-set limit.¹⁰⁵ The performance of W1–F12 theory has been recently evaluated against an even larger set of 200 very accurate atomization energies,¹⁰⁶ for this extended data set of atomization energies W1–F12 theory achieves an RMSD of $0.72 \text{ kcal mol}^{-1}$. The computational details of W3lite-F12 and W1–F12 theories have been specified and rationalized in great detail in refs.,^{101,103} see also ref.¹⁰⁴ for a recent review. We note that all the CCSD(T) energy calculations involved in the W3lite-F12 energies were done with the MOLPRO program package,^{89,90} whereas the

post-CCSD(T) calculations were carried out with the MRCC program.^{107,108}

RESULTS AND DISCUSSION

The relative energies (ΔE_0), activation energies (ΔE^\ddagger), reaction energies (ΔE_r), and NICS (1 Å) values calculated at different levels are given in Table 1. Schematic reaction profile diagrams connecting the reactants, **ptC1**, **ptC2**, and **ptC3** and their dissociative products via the relevant transition states are shown in Figures 2–4, respectively. The rotational and centrifugal distortion constants, inertial axis dipole moment components, and absolute dipole moments calculated at the CCSD(T)/cc-pVTZ level of theory for all ptC isomers are collected in Table 2. The optimal geometries of **ptC1**, **ptC2**,

Table 2. Rotational and Centrifugal Distortion Constants (in MHz), and Inertial Axis Dipole Moment Components (in Debye; \perp^r Representation) of ptC Isomers of C_7H_2 Calculated at the CCSD(T)/cc-pVTZ Level of Theory^a

constant	ptC1	ptC2	ptC3
A_e	5247.30	7319.09	8109.33
B_e	4710.27	2071.46	1957.82
C_e	2482.15	1614.52	1577.07
Δ_j	0.3423×10^{-3}	0.7399×10^{-3}	0.9046×10^{-3}
Δ_K	0.4666×10^{-3}	0.4814×10^{-1}	0.6579×10^{-1}
Δ_{JK}	0.4593×10^{-3}	-0.6880×10^{-2}	-0.1134×10^{-1}
δ_j	0.1292×10^{-3}	-0.2815×10^{-3}	-0.3153×10^{-3}
δ_K	0.4928×10^{-3}	-0.2258×10^{-4}	-0.2047×10^{-4}
μ_a	-5.84	0.12	-3.65
μ_b	-	-0.92	-4.20
$ \mu $	5.84	0.93	5.56

^aCentrifugal distortion constants are from A-reduced Hamiltonian for **ptC1** whereas they are estimated using the S-reduced Hamiltonian for **ptC2** and **ptC3** considering the fact that they are approaching close to the prolate limit.

and **ptC3** calculated at different levels are documented in Tables 3, 4, and 5, respectively. Energetic and spectroscopic properties of other low-lying isomers (1–9) are documented in our earlier works.^{59,60} For brevity the Cartesian coordinates of the optimized geometries, total electronic energies, ZPVEs, harmonic vibrational frequencies, and infrared intensities of all ptC isomers calculated at different levels are given in the Supporting Information.

Activation and Reaction Energies. For **ptC1**, two different transition states have been identified with respect to

breaking of the C–C single bond connected to the ptC atom (see Figure 2). The activation energy calculated for pathways **A** and **B** are 11.02 and 32.96 kcal mol⁻¹, respectively, at the W1–F12 level of theory. The activation energy for pathway **A** is in good agreement with the value of 9.08 kcal mol⁻¹ calculated at the B3LYP/6-311++G(2d,2p) level of theory by an earlier theoretical work.¹⁵ Nevertheless, our theoretical study differs from this earlier theoretical work with respect to dissociation pathway **B**, where it was concluded that it rearranges to **ptC1** itself. However, we find via IRC calculations that both the pathways lead to isomer **6**. For pathway **B**, we have noticed a ring-expansion (five-membered ring to six-membered ring) from the bicarbene like transition state (**ptC1-TS-B**), which once again leads to **6**. The calculated reaction energy of -5.65 kcal mol⁻¹ for **ptC1** indicates the exothermic nature of the minimum-energy pathway in arriving at **6**.

With respect to **ptC2** and **ptC3** dissociation pathways, altogether we have located seven different transition states (see Figures 3 and 4). Although it appears like four different transition states each, we note that pathway **C** for **ptC2** and pathway **D** for **ptC3** energetically lead to the same transition state. In both **ptC2** and **ptC3**, pathways **A** and **B** lead to isomer **2**. Isomer **3** is obtained via pathway **D** in **ptC2** and pathway **C** in **ptC3**. The lowest activation energy path (pathway **C** in **ptC2** and pathway **D** in **ptC3**) in both cases lead to a bicarbene structure, whose geometry is very similar to **ptC3** except that one of the C–C bonds connected to the ptC atom is broken. One can also arrive at **ptC2** geometry from this bicarbene structure by internal rotation of its C–C bond connected to the three-membered ring. It is worth to note that, in terms of relative energy, **ptC2** is more stable than the bicarbene product (isomer **BC**).¹⁰⁹ Therefore, for pathway **C** alone, the reaction energy is endothermic (3.27 kcal mol⁻¹). Though this pathway has the lowest activation energy (6.11 kcal mol⁻¹) compared to other pathways in **ptC2**, thermodynamically the reactant is more stable than the product. On the contrary, for the lowest activation energy path (pathway **D**) in **ptC3**, the reaction energy is near-zero at the W1–F12 level of theory. Without ZPVE correction, the reaction energy is still slightly exothermic (-0.22 kcal mol⁻¹; see Table S9 in the Supporting Information).

Considering the proximity of these three isomers (**ptC2**, **BC**, and **ptC3**), potential interconversions of **BC** to **ptC2** and **BC** to **ptC3** are quite likely. Moreover, whether it is **ptC2** or **ptC3**, the reaction energies for isomer **2** and **3** are highly exothermic though they are not arrived from the lowest activation energy pathways. Nevertheless, among these two products (**2** and **3**),

Table 3. Optimal Geometry Parameters (Å and Degrees) of Isomer ptC1 of C_7H_2 Calculated at Different Levels

parameter	cc-pVDZ		cc-pVTZ		6-311+G(d,p)	aug-cc-pVQZ
	MP2	CCSD(T)	MP2	CCSD(T)	B3LYP	MP2
$R(C_2C_4;C_3C_5)$	1.3836	1.3788	1.3652	1.3593	1.3490	1.3607
$R(C_1C_2;C_1C_3)$	1.4869	1.4858	1.4750	1.4727	1.4694	1.4728
$R(C_1C_4;C_1C_5)$	1.5175	1.5383	1.4965	1.5151	1.5077	1.4905
$R(C_2C_6;C_3C_7)$	1.4289	1.4438	1.4162	1.4303	1.4247	1.4142
$R(C_6C_7)$	1.3949	1.3907	1.3826	1.3772	1.3736	1.3802
$R(C_6H_8;C_7H_9)$	1.0941	1.0956	1.0804	1.0809	1.0821	1.0801
$\theta(C_2C_1C_3)$	94.94	95.71	94.62	95.32	95.61	94.64
$\theta(C_2C_1C_4;C_3C_1C_5)$	54.83	54.22	54.69	54.10	53.87	54.67
$\theta(C_1C_2C_6;C_1C_3C_7)$	116.35	115.80	116.58	116.10	115.81	116.56
$\theta(C_2C_6H_8;C_3C_7H_9)$	126.10	125.76	126.19	125.86	125.87	126.23

Table 4. Optimal Geometry Parameters (Å and Degrees) of Isomer ptC2 of C₇H₂ Calculated at Different Levels

parameter	cc-pVDZ		cc-pVTZ		6-311+G(d,p)	aug-cc-pVQZ
	MP2	CCSD(T)	MP2	CCSD(T)	B3LYP	MP2
R(C ₂ C ₄)	1.3521	1.3402	1.3325	1.3231	1.3123	1.3284
R(C ₃ C ₅)	1.3685	1.3921	1.3490	1.3609	1.3532	1.3449
R(C ₁ C ₂)	1.6112	1.7859	1.5903	1.6928	1.6950	1.5850
R(C ₁ C ₃)	1.4211	1.3972	1.4086	1.3917	1.3928	1.4064
R(C ₁ C ₄)	1.4210	1.4426	1.4081	1.4249	1.4219	1.4058
R(C ₁ C ₅)	1.5742	1.5282	1.5519	1.5271	1.5165	1.5456
R(C ₃ C ₇)	1.4084	1.4205	1.3937	1.4053	1.3937	1.3916
R(C ₇ C ₈)	1.2341	1.2306	1.2164	1.2120	1.2034	1.2142
R(C ₄ H ₆)	1.0892	1.0912	1.0748	1.0762	1.0779	1.0745
R(C ₈ H ₉)	1.0768	1.0799	1.0624	1.0642	1.0635	1.0624
θ(C ₄ C ₁ C ₂)	52.51	42.60	52.34	49.31	48.84	52.32
θ(C ₁ C ₄ C ₂)	70.99	79.75	70.88	75.95	76.51	70.79
θ(C ₁ C ₄ H ₆)	132.11	128.57	132.35	129.82	130.13	132.35
θ(C ₄ C ₁ C ₅)	128.99	139.81	129.49	135.64	134.99	129.83
θ(C ₁ C ₃ C ₇)	134.99	140.16	135.06	138.30	139.28	135.00
θ(C ₂ C ₁ C ₅)	178.50	172.59	178.17	175.06	176.18	177.85
θ(C ₄ C ₁ C ₃)	176.93	163.57	176.56	169.02	169.77	176.21
θ(C ₃ C ₇ C ₈)	179.66	179.48	179.51	179.82	179.67	179.00
θ(C ₇ C ₈ H ₉)	179.82	179.73	179.93	179.84	179.87	179.66

Table 5. Optimal Geometry Parameters (Å and Degrees) of Isomer ptC3 of C₇H₂ Calculated at Different Levels

parameter	cc-pVDZ		cc-pVTZ		6-311+G(d,p)	aug-cc-pVQZ
	MP2	CCSD(T)	MP2	CCSD(T)	B3LYP	MP2
R(C ₂ C ₄)	1.3527	1.3471	1.3327	1.3274	1.3153	1.3284
R(C ₃ C ₅)	1.3689	1.3770	1.3486	1.3531	1.3473	1.3442
R(C ₁ C ₂)	1.4341	1.4403	1.4237	1.4286	1.4260	1.4227
R(C ₁ C ₃)	1.4390	1.4183	1.4295	1.4102	1.4093	1.4282
R(C ₁ C ₄)	1.5913	1.6738	1.5644	1.6286	1.6447	1.5561
R(C ₁ C ₅)	1.5571	1.5714	1.5314	1.5491	1.5311	1.5245
R(C ₃ C ₇)	1.4067	1.4211	1.3917	1.4053	1.3931	1.3897
R(C ₇ C ₈)	1.2353	1.2312	1.2177	1.2127	1.2044	1.2157
R(C ₂ H ₆)	1.0904	1.0922	1.0760	1.0769	1.0784	1.0759
R(C ₈ H ₉)	1.0771	1.0800	1.0626	1.0643	1.0636	1.0627
θ(C ₅ C ₃ C ₁)	67.30	68.39	66.83	68.16	67.44	66.64
θ(C ₅ C ₃ C ₇)	158.18	155.10	158.67	155.72	155.09	158.94
θ(C ₃ C ₁ C ₂)	127.21	134.59	125.78	132.44	133.12	125.05
θ(C ₁ C ₂ H ₆)	133.25	131.70	133.46	132.07	132.24	133.42
θ(C ₄ C ₁ C ₂)	52.81	50.58	52.72	50.96	50.11	52.74
θ(C ₅ C ₁ C ₄)	125.78	120.27	127.45	122.43	122.42	128.17
θ(C ₂ C ₁ C ₅)	178.59	170.85	179.84	173.38	172.53	179.09
θ(C ₄ C ₁ C ₃)	179.98	174.83	178.49	176.60	176.77	177.79
θ(C ₃ C ₇ C ₈)	178.52	179.36	178.57	179.37	179.44	177.93
θ(C ₇ C ₈ H ₉)	178.68	179.14	178.88	179.28	179.20	178.61

the pathway leading to the experimentally known ring-chain isomer of C₇H₂ (**2**) is ~1.90 kcal mol⁻¹ more exothermic than the pathway leading to **3** at the W1-F12 level of theory. At the W3lite-F12 level of theory, these two isomers are energetically separated by 2.08 kcal mol⁻¹. We note that both kinetic and thermodynamic considerations govern the formation of a molecule in electrical discharges.⁷⁸ Although **2** alone was detected by FTMW spectroscopy⁵⁵ and **3** is yet to be observed,¹¹⁰ we hope that with a different choice of a precursor molecule and considering other kinetic factors such as collision time in the nozzle, the latter isomer could also be detected. In addition, it is worth mentioning here that **2** has an inertial axis dipole moment component in two directions (see Table 2), whereas for **3** it is in one direction.^{55,110,111} Therefore, both *a*-

and *b*-type rotational transitions are possible for **2** and only *b*-type rotational transition is possible for **3**. However, the latter type of transitions depend on the moment of inertia of both *A* and *C*. Therefore, a much more accurate theoretical prediction of its bond lengths and bond angles is required, which could be done in the future.

General Remarks on ptC Isomers in C₇H₂. The NICS (1 Å) values calculated at the B3LYP/6-311+G(d,p) level of theory for ptC1, ptC2, and ptC3 are -11.45, -7.72, and -9.02 ppm, respectively. The negative values indicate their aromatic nature, although they are not synthetically viable to date. We note that the reported NICS (1 Å) value for benzene molecule calculated at the same level of theory is -10.20 ppm.¹¹² Though the aromatic index of ptC1 is slightly higher

than benzene, it remains elusive to date. Nonetheless, we leave this discussion with a caveat that NICS is a response property whereas energetic stability is a ground-state property.¹¹³ The rotational constants (A_0 , B_0 , and C_0) obtained from the equilibrium geometry at the CCSD(T)/cc-pVTZ level of theory (see Table 2) reveal that all the ptC isomers studied here are asymmetric tops. The centrifugal distortion constants (Δ_p , Δ_K , Δ_{JK} , etc.) are also computed for these hypothetical molecules with a hope that they will assist the efforts of microwave spectroscopists. We note that the experimentally measured values are vibrationally averaged rotational constants (A_0 , B_0 , and C_0), whereas the values reported by us lacks the effects of corrections from the zero-point vibrational motion. Also, our calculations on these molecules lack the effect of core–valence correlation in their optimal geometries at the CCSD(T)/cc-pVTZ level of theory.¹¹⁴ Although these two effects may significantly improve the rotational constants, which would be helpful for the accurate simulation to search for the rotational transitions of any of the ptC isomers, such an endeavor is beyond the scope of our present work and would be undertaken in a forthcoming study. While the inertial axis dipole moment components are in two directions (μ_a and μ_b) for isomers **ptC2** and **ptC3**, it is in only one direction for **ptC1**. Therefore, for the latter, only one type of rotational transition is possible whereas for the former both *a*- and *b*-type rotational transitions are possible. On the basis of the absolute dipole moment values, one can say that **ptC1** and **ptC3** are more polar than **ptC2**. The total dipole moments of **ptC1**, **ptC2**, and **ptC3** are 5.84, 0.92, and 5.69 D, respectively, at the CCSD(T)/cc-pVTZ level of theory. The reason why **ptC2** is less polar could be attributed to the fact that the H atom and ethynyl ($-\text{C}\equiv\text{C}-\text{H}$) group are in opposite directions (unlike **ptC3**), and this geometrical arrangement affects the electron density distribution. Consequently, it cancels out the net dipole moment for **ptC2** to a large extent.

The optimal geometry parameters of **ptC1**, **ptC2**, and **ptC3** calculated at different levels are collected in Tables 3, 4, and 5, respectively. The atom numbering scheme we have adopted for the ptC isomers of C_7H_2 is given in Figure 5. At all levels of




Figure 5. Atom numbering scheme we have adopted for ptC isomers.

theory for all ptC isomers, the longer bond lengths of C_1C_2 , C_1C_3 , C_1C_4 , and C_1C_5 , and the double bond distance of C_2C_4 and C_3C_5 reflect their diradical valence structures given in Figure 1. The C_6C_7 bond length for **ptC1** at all levels is once again close to a double bond distance whereas the C_7C_8 bond length for **ptC2** and **ptC3** is that of a triple bond distance at all levels. While the C–H bond distance in **ptC1** is not surprising, the C–H bonds connected to the ethynyl group in **ptC2** and **ptC3** are somewhat shorter as they are connected to a *sp* hybridized carbon in both cases. Overall, with respect to cc-pVDZ basis set results, the bond lengths are found to be

systematically overestimated at the MP2 and CCSD(T) levels. These results are consistent with earlier observations and is largely due to the lack of higher angular momentum polarization functions.^{115–120} The near 180 deg bond angles of $\text{C}_3\text{C}_7\text{C}_8$ and $\text{C}_7\text{C}_8\text{H}_9$ at all levels in **ptC2** and **ptC3** confirm that the ethynyl chain is almost linear.

Since W1–F12 theory approximates the CCSD(T) basis-set-limit energy, it is of interest to estimate whether the contributions from post-CCSD(T) excitations are likely to be significant for the molecules considered in this work. The percentage of the total atomization energy (TAE) accounted for by the quasiperturbative triple excitations, %TAE[(T)],^{100,104,105,121,122} has been shown to be a reliable energy-based diagnostic for the importance of post-CCSD(T) contributions to the TAEs. It has been found that %TAE[(T)] values smaller than ~5% indicate that post-CCSD(T) contributions should normally not exceed 0.5 kcal mol⁻¹.¹⁰⁴ The %TAE[(T)] values for the C_7H_2 isomers considered in this work range between 2.8–3.7% (see Table S16 of the Supporting Information). These values suggest that the considered species are dominated by dynamical correlation effects and that post-CCSD(T) contributions to the TAEs should not exceed the 0.5 kcal mol⁻¹ mark. Thus, the predicted reaction energies and barrier heights obtained in the present work are expected to be little affected by post-CCSD(T) contributions.

CONCLUSIONS

Three isomers of C_7H_2 with a ptC atom and their dissociation pathways of C–C single bonds connected to the ptC atom have been theoretically studied. Energetically, isomer **ptC1** was found to lie ~5.78 kcal mol⁻¹ above the experimentally well-known and astronomically relevant cumulene carbene isomer of C_7H_2 (**9**) at the W3lite-F12 level of theory. Therefore, the dissociation of this molecule was studied in detail including IRC calculations to examine its kinetic stability. Key spectroscopic parameters relevant to microwave spectroscopists are also documented. We conclude that both the dissociation pathways for **ptC1** lead to **6**. However, the latter also remains elusive to date though thermodynamically more stable than **9**.

Though **ptC2** and **ptC3** can be considered as high-energy isomers from the thermodynamic perspective, their dissociation (except the lowest activation energy path) leads to low-energy isomers **2** and **3**, depending upon which C–C bond connected to the ptC atom is being broken. Among these two products, **2** is already identified by FTMW spectroscopy and **3** lies just 2.08 kcal mol⁻¹ above **2** at the W3lite-F12 level of theory. In all the cases (except pathway C in **ptC2**), the reaction appears to be exothermic. In conclusion, the dissociation pathways of ptC isomers lead to three low-energy isomers (**2**, **3**, and **6**) of C_7H_2 . However, the only low-lying ring–chain carbene identified to date is **2** and the other two isomers (**3** and **6**) remain elusive though both of them are energetically more stable than **9**.

In this work, by studying the dissociation pathways of ptC isomers, we have shown that the high-energy ptC isomers (**ptC2** and **ptC3**) serve as reactive intermediates to an experimentally known carbene molecule (**2**). Nevertheless, the most obvious question in front of us is, is it possible to identify the ptC isomers within the C_7H_2 elemental composition? On the basis of the activation energies and the reaction energies calculated in this work, we are hopeful that

the identification of the **ptC1** isomer is viable. If not, at least their dissociative products, isomers **3** and **6**, are the potential candidates for detection. We believe that our theoretical studies on these compounds would motivate the experimentalists in the synthesis and identification of these molecules in the future.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.jpca.8b08809](https://doi.org/10.1021/acs.jpca.8b08809).

Cartesian coordinates of the optimized geometries, total electronic energies, ZPVEs, harmonic vibrational frequencies, and infrared intensities of all ptC isomers calculated at different levels (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*(A.K.) E-mail: amir.karton@uwa.edu.au.

*(V.S.T.) E-mail: venkateshtsv@gmail.com.

ORCID

Krishnan Thirumoorthy: [0000-0003-0920-4547](https://orcid.org/0000-0003-0920-4547)

Amir Karton: [0000-0002-7981-508X](https://orcid.org/0000-0002-7981-508X)

Venkatesan S. Thimmakonda: [0000-0002-7505-077X](https://orcid.org/0000-0002-7505-077X)

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Initial part of this work is supported by a research grant (Project No: YSS/2015/000099) from the Science and Engineering Research Board, Department of Science and Technology (DST), New Delhi, Government of India (to VST). One of the authors (AK) thank Australian Research Council (ARC) Future Fellowship (Project No. FT170100373) for support received in part. We gratefully acknowledge the generous allocation of computing time from the National Computational Infrastructure (NCI) National Facility, and system administration support provided by the Faculty of Science at the University of Western Australia (UWA) to the Linux cluster of the Karton group. Additional computational facility provided (for KT) at the VIT University to carry out part of this work is also gratefully acknowledged.

■ REFERENCES

- (1) van't Hoff, J. H. A Suggestion Looking to the Extension Into Space of the Structural Formulas at Present Used in Chemistry. And A Note Upon the Relation Between the Optical Activity and the Chemical Constitution of Organic Compounds. *Arch. Neerl. Sci. Exactes Nat.* **1874**, *9*, 445–454.
- (2) Le-Bel, J. A. On the Relations Which Exist Between the Atomic Formulas of Organic Compounds and the Rotatory Power of Their Solutions. *Bull. Soc. Chim. Fr.* **1874**, *22*, 337.
- (3) Keese, R. Carbon Flatland: Planar Tetracoordinate Carbon and Fenestranes. *Chem. Rev.* **2006**, *106*, 4787–4808.
- (4) Li, X.; Wang, L. S.; Boldyrev, A. I.; Simons, J. Tetracoordinated Planar Carbon in the Al_4C^- Anion. A Combined Photoelectron Spectroscopy and Ab Initio Study. *J. Am. Chem. Soc.* **1999**, *121*, 6033–6038.
- (5) Li, X.; Zhang, H. F.; Wang, L. S.; Geske, G. D.; Boldyrev, A. I. Pentaatomic Tetracoordinate Planar Carbon, $[CAL_4]^{2-}$: A New Structural Unit and Its Salt Complexes. *Angew. Chem., Int. Ed.* **2000**, *39*, 3630–3633.

- (6) Wang, L. S.; Boldyrev, A. I.; Li, X.; Simons, J. Experimental Observation of Pentaatomic Tetracoordinate Planar Carbon-Containing Molecules. *J. Am. Chem. Soc.* **2000**, *122*, 7681–7687.
- (7) Li, X.; Zhai, H. J.; Wang, L. S. Photoelectron Spectroscopy of Pentaatomic Tetracoordinate Planar Carbon Molecules: CA_3Si and CA_3Ge . *Chem. Phys. Lett.* **2002**, *357*, 415–419.
- (8) Xu, J.; Zhang, X.; Yu, S.; Ding, Y.; Bowen, K. H. Identifying the Hydrogenated Planar Tetracoordinate Carbon: A Combined Experimental and Theoretical Study of CA_4H and CA_4H_2 . *J. Phys. Chem. Lett.* **2017**, *8*, 2263–2267.
- (9) Monkhorst, H. J. Activation Energy for Interconversion of Enantiomers Containing an Asymmetric Carbon Atom without Breaking Bonds. *Chem. Commun.* **1968**, 1111–1112.
- (10) Hoffmann, R.; Alder, R. W.; Wilcox, C. F., Jr. Planar Tetracoordinate Carbon. *J. Am. Chem. Soc.* **1970**, *92*, 4992–4993.
- (11) Collins, J. B.; Dill, J. D.; Jemmis, E. D.; Apeloig, Y.; Schleyer, P. v. R.; Seeger, R.; Pople, J. A. Stabilization of Planar Tetracoordinate Carbon. *J. Am. Chem. Soc.* **1976**, *98*, 5419–5427.
- (12) Sorger, K.; Schleyer, P. v. R. Planar and Inherently Non-Tetrahedral Tetracoordinate Carbon - a Status-Report. *J. Mol. Struct.: THEOCHEM* **1995**, *338*, 317–346.
- (13) Rottger, D.; Erker, G. Compounds Containing Planar-Tetracoordinate Carbon. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 812–827.
- (14) Siebert, W.; Gunale, A. Compounds Containing a Planar-Tetracoordinate Carbon Atom as Analogues of Planar Methane. *Chem. Soc. Rev.* **1999**, *28*, 367–371.
- (15) Perez, N.; Heine, T.; Barthel, R.; Seifert, G.; Vela, A.; Mendez-Rojas, M. A.; Merino, G. Planar Tetracoordinate Carbons in Cyclic Hydrocarbons. *Org. Lett.* **2005**, *7*, 1509–1512.
- (16) Merino, G.; Mendez-Rojas, M. A.; Vela, A.; Heine, T. Recent Advances in Planar Tetracoordinate Carbon Chemistry. *J. Comput. Chem.* **2007**, *28*, 362–372.
- (17) Priyakumar, U. D.; Sastry, G. N. A System with Three Contiguous Planar Tetracoordinate Carbons Is Viable: A Computational Study on a $C_6H_6^{2+}$ Isomer. *Tetrahedron Lett.* **2004**, *45*, 1515–1517.
- (18) McGrath, M. P.; Radom, L. Alkapanes: A Class of Neutral Hydrocarbons Containing a Potentially Planar Tetracoordinate Carbon. *J. Am. Chem. Soc.* **1993**, *115*, 3320–3321.
- (19) Lyons, J. E.; Rasmussen, D. R.; McGrath, M. P.; Nobes, R. H.; Radom, L. Octaplane: A Saturated Hydrocarbon with a Remarkably Low Ionization Energy Leading to a Cation with a Planar Tetracoordinate Carbon Atom. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1667–1668.
- (20) Radom, L.; Rasmussen, D. R. The Planar Carbon Story. *Pure Appl. Chem.* **1998**, *70*, 1977–1984.
- (21) Rasmussen, D. R.; Radom, L. Planar-Tetracoordinate Carbon in a Neutral Saturated Hydrocarbon: Theoretical Design and Characterization. *Angew. Chem., Int. Ed.* **1999**, *38*, 2875–2878.
- (22) Rasmussen, D. R.; Radom, L. Hemispiroalkapanes: Hydrocarbon Cage Systems with a Pyramidal-Tetracoordinate Carbon Atom and Remarkable Basicity. *Chem. - Eur. J.* **2000**, *6*, 2470–2483.
- (23) Pancharatna, P. D.; Mendez-Rojas, M. A.; Merino, G.; Vela, A.; Hoffmann, R. Planar Tetracoordinate Carbon in Extended Systems. *J. Am. Chem. Soc.* **2004**, *126*, 15309–15315.
- (24) Wang, Z. X.; Zhang, C. G.; Chen, Z. F.; Schleyer, P. v. R. Planar Tetracoordinate Carbon Species Involving Beryllium Substituents. *Inorg. Chim. Acta.* **2008**, *47*, 1332–1336.
- (25) Crigger, C.; Wittmaack, B. K.; Tawfik, M.; Merino, G.; Donald, K. J. Plane and Simple: Planar Tetracoordinate Carbon Centers in Small Molecules. *Phys. Chem. Chem. Phys.* **2012**, *14*, 14775–14783.
- (26) Menzel, M.; Steiner, D.; Winkler, H.-J.; Schweikart, D.; Mehle, S.; Fau, S.; Frenking, G.; Massa, W.; Berndt, A. Compounds with Planar Tetracoordinate Boron Atoms: Anti van't Hoff/Le Bel Geometries without Metal Centers. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 327–329.

- (27) Priyakumar, U. D.; Reddy, A. S.; Sastry, G. N. The Design of Molecules Containing Planar Tetracoordinate Carbon. *Tetrahedron Lett.* **2004**, *45*, 2495–2498.
- (28) Sateesh, B.; Reddy, A. S.; Sastry, G. N. Towards Design of the Smallest Planar Tetracoordinate Carbon and Boron Systems. *J. Comput. Chem.* **2007**, *28*, 335–343.
- (29) Cui, Z.-H.; Contreras, M.; Ding, Y.-H.; Merino, G. Planar Tetracoordinate Carbon versus Planar Tetracoordinate Boron: The Case of CB_4 and Its Cation. *J. Am. Chem. Soc.* **2011**, *133*, 13228–13231.
- (30) Yang, S.; Zhang, C. Theoretical Investigation of Obtaining Compounds with Planar Tetracoordinate Carbons by Frustrated Lewis Pairs. *J. Phys. Chem. A* **2015**, *119*, 8950–8957.
- (31) Zhang, C.; Jia, W.; Cao, Z. Stability Rules of Main-Group Element Compounds with Planar Tetracoordinate Carbons. *J. Phys. Chem. A* **2010**, *114*, 7960–7966.
- (32) Yanez, O.; Vasquez-Espinal, A.; Pino-Rios, R.; Ferraro, F.; Pan, S.; Osorio, E.; Merino, G.; Tiznado, W. Exploiting Electronic Strategies to Stabilize A Planar Tetracoordinate Carbon in Cyclic Aromatic Hydrocarbons. *Chem. Commun.* **2017**, *53*, 12112–12115.
- (33) Driess, M.; Aust, J.; Merz, K.; van Wüllen, C. van't Hoff-Le Bel Stranger: Formation of a Phosphonium Cation with a Planar Tetracoordinate Phosphorus Atom. *Angew. Chem., Int. Ed.* **1999**, *38*, 3677–3680.
- (34) Stephan, D. W. “Breaking the Rules”: A Planar Phosphonium Cation. *Angew. Chem., Int. Ed.* **2000**, *39*, 501–502.
- (35) Roy, D.; Corminboeuf, C.; Wannere, C. S.; King, R. B.; Schleyer, P. v. R. Planar Tetracoordinate Carbon Atoms Centered in Bare Four-membered Rings of Late Transition Metals. *Inorg. Chem.* **2006**, *45*, 8902–8906.
- (36) Wang, Z. X.; Schleyer, P. v. R. Construction Principles of “Hyparenes”: Families of Molecules with Planar Pentacoordinate Carbons. *Science* **2001**, *292*, 2465–2469.
- (37) Li, S. D.; Miao, C. Q.; Ren, G. M. D_{5h} Cu_5H_5X : Pentagonal Hydrocopper Cu_5H_5 Containing Pentacoordinate Planar Nonmetal Centers ($X = B, C, N, O$). *Eur. J. Inorg. Chem.* **2004**, *2004*, 2232–2234.
- (38) Erhardt, S.; Frenking, G.; Chen, Z.; Schleyer, P. v. R. Aromatic Boron Wheels with More than One Carbon Atom in the Center: C_2B_8 , $C_3B_9^+$, and $C_5B_{11}^+$. *Angew. Chem., Int. Ed.* **2005**, *44*, 1078–1082.
- (39) Grande-Aztatzi, R.; Cabellos, J. L.; Islas, R.; Infante, I.; Mercero, J. M.; Restrepo, A.; Merino, G. Planar Pentacoordinate Carbons in CB_5^{4-} Derivatives. *Phys. Chem. Chem. Phys.* **2015**, *17*, 4620–4624.
- (40) Vassilev-Galindo, V.; Pan, S.; Donald, K. J.; Merino, G. Planar Pentacoordinate Carbons. *Nat. Chem. Rev.* **2018**, *2*, 0114.
- (41) Ravell, E.; Jalife, S.; Barroso, J.; Orozco-Ic, M.; Hernandez-Juarez, G.; Ortiz-Chi, F.; Pan, S.; Cabellos, J. L.; Merino, G. Structure and Bonding in CE_5^- ($E = Al-Tl$) Clusters: Planar Tetracoordinate Carbon vs Pentacoordinate Carbon. *Chem. - Asian J.* **2018**, *13*, 1467–1473.
- (42) Pan, S.; Cabellos, J. L.; Orozco-Ic, M.; Chattaraj, P. K.; Zhao, L.; Merino, G. Planar Pentacoordinate Carbon in CGa^{5+} Derivatives. *Phys. Chem. Chem. Phys.* **2018**, *20*, 12350–12355.
- (43) Wang, Y.; Li, F.; Li, Y.; Chen, Z. Semi-Metallic Be_5C_2 Monolayer Global Minimum with Quasi-Planar Pentacoordinate Carbons and Negative Poisson's Ratio. *Nat. Commun.* **2016**, *7*, 11488.
- (44) Exner, K.; Schleyer, P. v. R. Planar Hexacoordinate Carbon: A Viable Possibility. *Science* **2000**, *290*, 1937–1940.
- (45) Ito, K.; Chen, Z. F.; Corminboeuf, C.; Wannere, C. S.; Zhang, X. H.; Li, Q. S.; Schleyer, P. v. R. Myriad Planar Hexacoordinate Carbon Molecules Inviting Synthesis. *J. Am. Chem. Soc.* **2007**, *129*, 1510–1511.
- (46) Zhang, C.-F.; Han, S.-J.; Wu, Y.-B.; Lu, H.-G.; Lu, G. Thermodynamic Stability versus Kinetic Stability: Is the Planar Hexacoordinate Carbon Species D_{3h} $CN_3Mg_3^+$ Viable? *J. Phys. Chem. A* **2014**, *118*, 3319–3325.
- (47) Averkiev, B. B.; Zubarev, D. Y.; Wang, L. M.; Huang, W.; Wang, L. S.; Boldyrev, A. I. Carbon Avoids Hypercoordination in CB_6 , CB_6^- , and $C_2B_5^-$ Planar Carbon-Boron Clusters. *J. Am. Chem. Soc.* **2008**, *130*, 9248–9249.
- (48) Forgy, C. C.; Schlimgen, A. W.; Mazziotti, D. Effects of Nitrogenous Substituent Groups on the Benzene Dication. *Mol. Phys.* **2018**, *116*, 1364–1368.
- (49) Wang, L. M.; Huang, W.; Averkiev, B. B.; Boldyrev, A. I.; Wang, L. S. CB_7^- : Experimental and Theoretical Evidence against Hypercoordinate Planar Carbon. *Angew. Chem., Int. Ed.* **2007**, *46*, 4550–4553.
- (50) Wang, G.; Rahman, A. K. F.; Wang, B. Ab Initio Calculations of Ionic Hydrocarbon Compounds with Heptacoordinate Carbon. *J. Mol. Model.* **2018**, *24*, 116.
- (51) Yang, L.-M.; Ganz, E.; Chen, Z.; Wang, Z.-X.; Schleyer, P. v. R. Four Decades of the Chemistry of Planar Hypercoordinate Compounds. *Angew. Chem., Int. Ed.* **2015**, *54*, 9468–9501.
- (52) Considering the diradical nature of these molecules, assigning proper IUPAC names is cumbersome. Therefore, they are named as didehydro derivatives. We note that the parent molecule names (C_7H_4) are indeed IUPAC names.
- (53) Apponi, A. P.; McCarthy, M. C.; Gottlieb, C. A.; Thaddeus, P. Laboratory Detection of Four New Cumulene Carbenes: H_2C_7 , H_2C_8 , H_2C_9 , and D_2C_{10} . *Astrophys. J.* **2000**, *530*, 357–361.
- (54) Dua, S.; Blanksby, S. J.; Bowie, J. H. Formation of Neutral C_7H_2 Isomers from Four Isometric C_7H_2 Radical Anion Precursors in the Gas Phase. *J. Phys. Chem. A* **2000**, *104*, 77–85.
- (55) McCarthy, M. C.; Travers, M. J.; Gottlieb, C. A.; Thaddeus, P. Laboratory Detection of the Ring-Chain Molecule C_7H_2 . *Astrophys. J.* **1997**, *483*, L139–L142.
- (56) Aoki, K.; Ikuta, S. The Singlet with a C_3 Ring: The Probable Candidate of HC_6N and C_7H_2 . *J. Mol. Struct.: THEOCHEM* **1994**, *310*, 229–238.
- (57) Alkorta, I.; Elguero, J. Polyynes vs. Cumulenes: Their Possible Use as Molecular Wires. *Struct. Chem.* **2005**, *16*, 77–79.
- (58) Sun, B. J.; Huang, C. H.; Tsai, M. F.; Sun, H. L.; Gao, L. G.; Wang, Y. S.; Yeh, Y. Y.; Shih, Y. H.; Sia, Z. F.; Chen, P. H.; et al. Synthesis of Interstellar 1,3,5-heptatriynylidyne, $C_7H(X^2\Pi)$, Via The Neutral-Neutral Reaction of Ground State Carbon Atom, $C(^3P)$, With Triacetylene, $HC_6H(X^1\Sigma_g^+)$. *J. Chem. Phys.* **2009**, *131*, 104305–13.
- (59) Thimmakondur, V. S. The Equilibrium Geometries of Heptatriynylidene, Cyclohepta-1,2,3,4-tetraen-6-yne, and Heptahexaenylidene. *Comput. Theor. Chem.* **2016**, *1079*, 1–10.
- (60) Thimmakondur, V. S.; Karton, A. Energetic and Spectroscopic Properties of the Low-Lying C_7H_2 Isomers: A High-Level Ab Initio Perspective. *Phys. Chem. Chem. Phys.* **2017**, *19*, 17685–17697.
- (61) Fulara, J.; Freivogel, P.; Forney, D.; Maier, J. P. Electronic Absorption Spectra of Linear Carbon Chains in Neon Matrices. III $HC_{2n+1}H$. *J. Chem. Phys.* **1995**, *103*, 8805–8810.
- (62) Ball, C. D.; McCarthy, M. C.; Thaddeus, P. Laser Spectroscopy of the Carbon Chains HC_7H and HC_9H . *Astrophys. J.* **1999**, *523*, L89–L91.
- (63) Ball, C. D.; McCarthy, M. C.; Thaddeus, P. Cavity Ringdown Spectroscopy of the Linear Carbon Chains HC_7H , HC_9H , $HC_{11}H$, and $HC_{13}H$. *J. Chem. Phys.* **2000**, *112*, 10149–10155.
- (64) Thaddeus, P.; Vrtilek, J. M.; Gottlieb, C. A. Laboratory and Astronomical Identification of Cyclopropenylidene, C_3H_2 . *Astrophys. J.* **1985**, *299*, L63–L66.
- (65) Cernicharo, J.; Gottlieb, C. A.; Guelin, M.; Killian, T. C.; Paubert, G.; Thaddeus, P.; Vrtilek, J. M. Astronomical Detection of H_2CCC . *Astrophys. J.* **1991**, *368*, L39–L41.
- (66) Cernicharo, J.; Gottlieb, C. A.; Guelin, M.; Killian, T. C.; Thaddeus, P.; Vrtilek, J. M. Astronomical Detection of H_2CCCC . *Astrophys. J.* **1991**, *368*, L43–L45.
- (67) Langer, W. D.; Velusamy, T.; Kuiper, T. B. H.; Peng, R.; McCarthy, M. C.; Travers, M. J.; Kovács, A.; Gottlieb, C. A.; Thaddeus, P. First Astronomical Detection of the Cumulene Carbon Chain Molecule H_2C_6 in TMC-1. *Astrophys. J.* **1997**, *480*, L63–L66.
- (68) Vrtilek, J. M.; Gottlieb, C. A.; Gottlieb, E. W.; Killian, T. C.; Thaddeus, P. Laboratory Detection of Propadienylidene, H_2CCC . *Astrophys. J.* **1990**, *364*, L53–L56.

- (69) Killian, T. C.; Vrtilek, J. M.; Gottlieb, C. A.; Gottlieb, E. W.; Thaddeus, P. Laboratory Detection of a Second Carbon Chain Carbene: Butatrienylidene, H_2CCCC . *Astrophys. J.* **1990**, *365*, L89–L92.
- (70) McCarthy, M. C.; Travers, M. J.; Kovács, A.; Chen, W.; Novick, S. E.; Gottlieb, C. A.; Thaddeus, P. Detection and Characterization of the Cumulene Carbenes H_2C_5 and H_2C_6 . *Science* **1997**, *275*, 518–520.
- (71) Spezzano, S.; Brünken, S.; Schilke, P.; Caselli, P.; Menten, K. M.; McCarthy, M. C.; Bizzocchi, L.; Treviño-Morales, S. P.; Aikawa, Y.; Schlemmer, S. Interstellar Detection of $c\text{-C}_3\text{D}_2$. *Astrophys. J., Lett.* **2013**, *769*, L19.
- (72) Kawaguchi, K.; Kasai, Y.; Ishikawa, S.; Kaifu, N. A Spectral-Line Survey Observation of IRC+10216 between 28 and 50 GHz. *Pub. Ast. Soc. Japan* **1995**, *43*, 853–876.
- (73) Bell, M. B.; Avery, L. W.; Watson, J. K. G. A Spectral-Line Survey of W51 from 17.6 to 22.0 GHz. *Astrophys. J., Suppl. Ser.* **1993**, *86*, 211–233.
- (74) Herbst, E.; Leung, C. M. Gas-Phase Production of Complex Hydrocarbons, Cyanopolynes, and Related Compounds in Dense Interstellar Clouds. *Astrophys. J., Suppl. Ser.* **1989**, *69*, 271–300.
- (75) Winnewisser, G.; Walmsley, C. M. The Detection of HC_3N and HC_7N in IRC+10216. *Astron. Astrophys.* **1978**, *70*, L37–L39.
- (76) Pardo, J. R.; Cernicharo, J.; Goicoechea, J. R. Observational Evidence of the Formation of Cyanopolynes in CRL 618 Through the Polymerization of HCN. *Astrophys. J.* **2005**, *628*, 275–282.
- (77) Bell, M. B.; Matthews, H. E. Detection of C_3N in the Spiral Arm Gas Clouds in the Direction of Cassiopeia A. *Astrophys. J.* **1995**, *438*, 223–225.
- (78) Travers, M. J.; McCarthy, M. C.; Gottlieb, C. A.; Thaddeus, P. Laboratory Detection of the Ring-Chain Molecule C_5H_2 . *Astrophys. J.* **1997**, *483*, L135–L138.
- (79) Bühl, M.; van Willen, C. Computational Evidence for a New C_{84} Isomer. *Chem. Phys. Lett.* **1995**, *247*, 63–68.
- (80) Schleyer, P. v. R.; Maerker, C.; Dransfeld, A.; Jiao, H.; van Eikema Hommes, N. J. R. Nucleus-Independent Chemical Shifts: A Simple and Efficient Aromaticity Probe. *J. Am. Chem. Soc.* **1996**, *118*, 6317–6318.
- (81) Möller, C.; Plesset, M. S. Note On An Approximation Treatment for Many-Electron Systems. *Phys. Rev.* **1934**, *46*, 618–622.
- (82) Purvis, G. D., III; Bartlett, R. J. A Full Coupled-Cluster Singles and Doubles Model: The Inclusion of Disconnected Triples. *J. Chem. Phys.* **1982**, *76*, 1910.
- (83) Stanton, J. F.; Gauss, J.; Watts, J. D.; Bartlett, R. J. A Direct Product Decomposition Approach for Symmetry Exploitation in Many-Body Methods. I. Energy calculations. *J. Chem. Phys.* **1991**, *94*, 4334.
- (84) Raghavachari, K.; Trucks, G. W.; Pople, J. A.; Head-Gordon, M. A Fifth-Order Perturbation Comparison of Electron Correlation Theories. *Chem. Phys. Lett.* **1989**, *157*, 479–483.
- (85) Bartlett, R. J.; Watts, J. D.; Kucharski, S. A.; Noga, J. Non-Iterative Fifth-Order Triple and Quadruple Excitation Energy Corrections In Correlated Methods. *Chem. Phys. Lett.* **1990**, *165*, 513–522.
- (86) Stanton, J. F. Why CCSD(T) Works: A Different Perspective. *Chem. Phys. Lett.* **1997**, *281*, 130–134.
- (87) Dunning, T. H., Jr. Gaussian Basis Sets for Use in Correlated Molecular Calculations. I. The Atoms Boron Through Neon and Hydrogen. *J. Chem. Phys.* **1989**, *90*, 1007.
- (88) CFOUR, *Coupled-Cluster Techniques for Computational Chemistry, A Quantum-Chemical Program Package* by Stanton, J. F.; Gauss, J.; Harding, M. E.; Szalay, P. G.; with contributions from Auer, A. A.; Bartlett, R. J.; Benedikt, U.; Berger, C.; Bernholdt, D. E.; Bomble, Y. J.; et al. For detailed information, see: <http://www.cfour.de>.
- (89) Werner, H.-J.; Knowles, P. J.; Knizia, G.; Manby, F. R.; Schütz, M.; Celani, P.; Györfy, W.; Kats, D.; Korona, T.; Lindh, R., et al. *MOLPRO A Package of Ab Initio Programs: MOLPRO 2012.1*; University College Cardiff Consultants Limited: Cardiff, U.K., 2012. See also: <http://www.molpro.net>.
- (90) Werner, H.-J.; Knowles, P. J.; Knizia, G.; Manby, F. R.; Schütz, M. Molpro: A General-Purpose Quantum Chemistry Program Package. *WIREs Comput. Mol. Sci.* **2012**, *2*, 242–253.
- (91) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti Correlation-Energy Formula Into a Functional of the Electron Density. *Phys. Rev. B: Condens. Matter Mater. Phys.* **1988**, *37*, 785.
- (92) Becke, A. D. Density-Functional Exchange-Energy Approximation with Correct Asymptotic Behavior. *Phys. Rev. A: At., Mol., Opt. Phys.* **1988**, *38*, 3098.
- (93) Krishnan, R.; Binkley, J. S.; Seeger, R.; Pople, J. A. Self-Consistent Molecular Orbital Methods. XX. A Basis Set for Correlated Wave Functions. *J. Chem. Phys.* **1980**, *72*, 650.
- (94) Clark, T.; Chandrasekhar, J.; Spitznagel, G. W.; Schleyer, P. v. R. Efficient Diffuse Function-Augmented Basis Sets for Anion Calculations. III. The 3-21+G Basis Set for First-Row Elements, Li–F. *J. Comput. Chem.* **1983**, *4*, 294–301.
- (95) Fukui, K. The Path of Chemical-Reactions - The IRC Approach. *Acc. Chem. Res.* **1981**, *14*, 363–368.
- (96) Hratchian, H. P.; Schlegel, H. B. Finding minima, transition states, and following reaction pathways on ab initio potential energy surfaces. *Theory and Applications of Computational Chemistry: The First 40 Years* **2005**, 195–249.
- (97) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A. et al. *Gaussian 09, Revision E.01*; Gaussian, Inc.: Wallingford, CT, 2013.
- (98) Gauss, J.; Stanton, J. F. Analytic CCSD(T) Second Derivatives. *Chem. Phys. Lett.* **1997**, *276*, 70–77.
- (99) Kesharwani, M. K.; Brauer, B.; Martin, J. M. L. Frequency and Zero-Point Vibrational Energy Scale Factors for Double-Hybrid Density Functionals (and Other Selected Methods): Can Anharmonic Force Fields Be Avoided? *J. Phys. Chem. A* **2015**, *119*, 1701–1714.
- (100) Karton, A.; Rabinovich, E.; Martin, J. M. L.; Ruscic, B. W4 Theory for Computational Thermochemistry: In Pursuit of Confident sub-kJ/mol Predictions. *J. Chem. Phys.* **2006**, *125*, 144108.
- (101) Karton, A.; Kaminker, I.; Martin, J. M. L. Economical Post-CCSD(T) Computational Thermochemistry Protocol and Applications to Some Aromatic Compounds. *J. Phys. Chem. A* **2009**, *113*, 7610.
- (102) Sylvetsky, N.; Peterson, K. A.; Karton, A.; Martin, J. M. L. Toward a W4-F12 Approach: Can Explicitly Correlated and Orbital-Based Ab Initio CCSD(T) Limits be Reconciled? *J. Chem. Phys.* **2016**, *144*, 214101.
- (103) Karton, A.; Martin, J. M. L. Explicitly Correlated Wn Theory: W1-F12 and W2-F12. *J. Chem. Phys.* **2012**, *136*, 124114.
- (104) Karton, A. A Computational Chemist's Guide to Accurate Thermochemistry for Organic Molecules. *WIREs Comput. Mol. Sci.* **2016**, *6*, 292–310.
- (105) Karton, A.; Daon, S.; Martin, J. M. L. W4–11: A High-Confidence Benchmark Dataset for Computational Thermochemistry Derived from First-Principles W4 Data. *Chem. Phys. Lett.* **2011**, *510*, 165–178.
- (106) Karton, A.; Sylvetsky, N.; Martin, J. M. L. W4–17: A Diverse and High-Confidence Dataset of Atomization Energies for Benchmarking High-Level Electronic Structure Methods. *J. Comput. Chem.* **2017**, *38*, 2063–2075.
- (107) Kállay, M.; Rolik, Z.; Csontos, J.; Nagy, P.; Samu, G.; Mester, D.; Csóka, J.; Szab, B.; Ladjanszki, I.; Szegedy, L., et al. *MRCC, A Quantum Chemical Program Suite*. For the current version, see: <http://www.mrcc.hu>.
- (108) Rolik, Z.; Szegedy, L.; Ladjanszki, I.; Ladoczki, B.; Kállay, M. An Efficient Linear-Scaling CCSD(T) Method Based on Local Natural Orbitals. *J. Chem. Phys.* **2013**, *139*, 094105.
- (109) We note that at the CCSD(T)/cc-pVTZ level of theory including ZPVE correction at the same level and a scaling factor of 0.9868, isomer BC is 2.73 kcal mol⁻¹ above ptC2 but 0.58 kcal mol⁻¹ below ptC3.

(110) Chandra, S.; Chang, Ch.; Musrif, P. G.; Patzer, A. B. C.; Kegel, W. H.; Sedlmayr, E. Suggestion for A Search of the Cyclic Molecule $c-C_7H_2$ in Cool Cosmic Objects. *Rom. Journ. Phys.* **2007**, *52*, 459–466.

(111) McCarthy, M. C.; Grabow, J.-U.; Travers, M. J.; Chen, W.; Gottlieb, C. A.; Thaddeus, P. Laboratory Detection of the Ring-Chain Carbenes HC_4N and HC_6N . *Astrophys. J.* **1999**, *513*, 305–310.

(112) Chen, Z.; Wannere, C. S.; Corminboeuf, C.; Puchta, R.; Schleyer, P. v. R. Nucleus-Independent Chemical Shifts (NICS) as an Aromaticity Criterion. *Chem. Rev.* **2005**, *105*, 3842–3888.

(113) Foroutan-Nejad, C. Interatomic Magnetizability: A QTAIM-Based Approach toward Deciphering Magnetic Aromaticity. *J. Phys. Chem. A* **2011**, *115*, 12555–12560.

(114) Peterson, K. A.; Dunning, T. H., Jr Accurate Correlation Consistent Basis Sets for Molecular Core-Valence Correlation Effects: The Second Row Atoms Al–Ar, and the First Row Atoms B–Ne Revisited. *J. Chem. Phys.* **2002**, *117*, 10548.

(115) Spackman, P. R.; Jayatilaka, D.; Karton, A. Basis Set Convergence of CCSD(T) Equilibrium Geometries Using a Large and Diverse Set of Molecular Structures. *J. Chem. Phys.* **2016**, *145*, 104101.

(116) Feller, D.; Peterson, K. A. Probing the Limits of Accuracy in Electronic Structure Calculations: Is Theory Capable of Results Uniformly Better Than “Chemical Accuracy”? *J. Chem. Phys.* **2007**, *126*, 114105.

(117) Wang, S.; Schaefer, H. F., III The Small Planarization Barriers for the Amino Group in the Nucleic Acid Bases. *J. Chem. Phys.* **2006**, *124*, 044303.

(118) Bak, K. L.; Gauss, J.; Jørgensen, P.; Olsen, J.; Helgaker, T.; Stanton, J. F. The Accurate Determination of Molecular Equilibrium Structures. *J. Chem. Phys.* **2001**, *114*, 6548.

(119) Martin, J. M. L.; Taylor, P. R. Structure and Vibrations of Small Carbon Clusters from Coupled-Cluster Calculations. *J. Phys. Chem.* **1996**, *100*, 6047–6056.

(120) Xie, Y.; Scuseria, G. E.; Yates, B. F.; Yamaguchi, Y.; Schaefer, H. F. Methylnitrene: Theoretical Predictions of its Molecular Structure and Comparison with the Conventional C–N Single Bond in Methylamine. *J. Am. Chem. Soc.* **1989**, *111*, 5181–5185.

(121) Karton, A.; Martin, J. M. L. Performance of W4 Theory for Spectroscopic Constants and Electrical Properties of Small Molecules. *J. Chem. Phys.* **2010**, *133*, 144102.

(122) Fogueri, U. R.; Kozuch, S.; Karton, A.; Martin, J. M. L. A Simple DFT-Based Diagnostic for Nondynamical Correlation. *Theor. Chem. Acc.* **2013**, *132*, 1291.

(123) We note that the relative energy differences for isomers **1–9** and also for the three ptC isomers (**ptC1**, **ptC2**, and **ptC3**) were calculated at the W3lite-F12 level of theory. For isomers **10**, **11**, and **15**, the relative energies mentioned in [Figure 1](#) were estimated at the fc-CCSD(T)/cc-pVTZ level of theory (ZPVE corrected and scaled). As far as isomer **16** is concerned, our calculations are not converged. Therefore, the relative energy difference reported here for this particular isomer is taken from [ref 54](#), estimated at the B3LYP/aug-cc-pVDZ//B3LYP/6-31G(d) level of theory